SUPPLEMENTARY MATERIAL

Evaluation of the renal function among sickle cell patients: a cross-sectional study

Talal Qadah, 1,2 Anwar Refaei 1,2

¹Department of Medical Laboratory Sciences, Faculty of Applied Medical Sciences, King Abdulaziz University, Jeddah; ²Hematology Research Unit, King Fahad Medical Research Center, King Abdulaziz University, Jeddah, Saudi Arabia

Correspondence: Talal Qadah, Department of Medical Laboratory Sciences, Faculty of Applied Medical Sciences, King Abdulaziz University, 21589 Jeddah, Saudi Arabia.

Tel.: +966 (0) 55 554 0081. E-mail: thqadah@kau.edu.sa

Key words: sickle cell disorders, hemoglobinopathy, renal, creatinine, urea.

Supplementary Table 1. The inclusion and exclusion criteria.

Inclusion criteria	Exclusion criteria				
Patients from Saudi Arabia	Patient's residence in Saudi Arabia but not citizen				
All genders	Pregnant women				
Availability	Sickle pain crises and urinary tract infection				

Supplementary Table 2. Results of hemoglobin electrophoresis among cases with sickle cell anemia.

Hemoglobin type	Total (n=242)	Children (n=120)	Adults (n=122)	p
HbS	67.4±23.8	69.99±22.01	64.84±22.01	0.092
HbA ₁	22.4±27.3	18.00±25.41	26.76±28.42	0.012
HbF	6.3±5.5	8.12±5.98	4.50±4.29	< 0.001
HbA ₂	3.9±0.9	3.88 ± 0.85	3.94±1.01	0.642

Supplementary Table 3. Blood picture of participant male control subjects and male cases with sickle cell disorder.

Hemoglobin type	Control (n=52)		Cases (n=115)		
Blood picture	Mean	SD	Mean	SD	р
White blood cell count (×10 ³ /μL)	7.26	1.61	13.02	5.87	< 0.001
Red blood cell count (×10 ⁶ /μL)	5.16	0.60	3.70	1.10	< 0.001
Hemoglobin (g/dL)	12.62	1.65	9.20	2.09	< 0.001
Hematocrit (%)	40.02	4.02	28.01	6.96	< 0.001
Mean corpuscular volume (fL)	77.94	7.14	77.13	10.00	0.630
Mean corpuscular hemoglobin (pg)	24.61	3.32	25.40	3.76	0.232
MCHC (g/dL)	31.44	1.89	33.05	2.27	< 0.001
Red cell distribution width (%)	14.11	1.89	19.31	4.00	< 0.001
Platelets (×10 ⁶ /μL)	304.36	86.76	408.11	173.32	< 0.001

SD, standard deviation; MCHC, mean corpuscular hemoglobin concentration.

Supplementary Table 4. Serum electrolytes (sodium and potassium) levels of participant control subjects and cases with sickle cell anemia (Hb SS) compared with those with Hb AS.

-	Control (n=91)		Cases (n=242)		n
Electrolytes	Mean	SD	Mean	SD	р
Sodium (mEq/L)	136.68	2.40	137.43	3.27	0.046
Potassium (mEq/L)	3.97	0.49	4.14	0.52	0.009
	Hb SS (n=172)		Hb AS (n=70)		n
Electrolytes	Mean	SD	Mean	SD	р
Sodium (mEq/L)	137.18	3.26	138.06	3.24	0.059
Potassium (mEq/L)	4.12	0.54	4.19	0.46	0.339

SD, standard deviation.

Supplementary Table 5. Serum electrolytes (sodium and potassium) levels of participant control subjects and Hb SS cases with sickle cell anemia as well as control subjects and Hb AS cases with sickle cell trait.

	Control (n=91)		Hb SS Cases (n=172)		n
Electrolytes	Mean	SD	Mean	SD	р
Sodium (mEq/L)	136.68	2.40	137.18	3.26	0.200
Potassium (mEq/L)	3.97	0.49	4.12	0.54	0.034
	Control (n=91)		Hb AS Cases (n=70)		
Electrolytes	Mean	SD	Mean	SD	р
Sodium (mEq/L)	136.68	2.40	138.06	3.24	0.002
Potassium (mEq/L)	3.97	0.49	4.19	0.46	0.005

SD, standard deviation.

