



eISSN 1877-9352

Italian Journal of Medicine

<https://www.italjmed.org/ijm>

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The Early Access service lets users access peer-reviewed articles well before print/regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Italian Journal of Medicine** is, therefore, E-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination, and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear in a regular issue of the journal.

The E-publishing of this PDF file has been approved by the authors.

Please cite this article as:

Faggian G, Faggian R, Salzano M, et al. **Right ventricle migration of a peripherally inserted central catheter: case report and literature review.** *Ital J Med* doi: 10.4081/itjm.2025.2123

Submitted: 24-07-2025

Accepted: 02-08-2025

 © the Author(s), 2025
Licensee PAGEPress, Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.
All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Right ventricle migration of a peripherally inserted central catheter: case report and literature review

Guido Faggian,¹ Roberto Faggian,² Michela Salzano,³ Ciro Stavolo,¹
Teresa Argenziano,⁴ Andrea Diglio,⁵ Angela Faggian⁵

¹Department of Diagnostic Imaging, San Felice a Canello Hospital, Maddaloni (CE); ²CEDIAL Dialysis Center, San Cipriano d'Aversa (CE); ³Local Health Authority 1 Center, Naples; ⁴Department of Neuroscience, Reproductive Science and Dentistry, University of Naples Federico II; ⁵Department of Diagnostic Imaging, National Hospital Healthcare Organization "San Pio", Benevento, Italy

Correspondence: Guido Faggian, Department of Diagnostic Imaging, San Felice a Canello Hospital, Maddaloni (CE), Italy.
E-mail: guidofaggian@libero.it

Key words: PICC fragmentation, endovascular retrieval, diagnostic imaging, intravascular foreign body, patient safety.

Contributions: the authors received no contributions from any individual or organization in relation to this work.

Conflict of interest: the authors declare no competing interests. All authors have reviewed the manuscript and confirm the accuracy of the data and the integrity of the work.

Ethics approval and consent to participate: the ethics committee deemed formal approval unnecessary, as no experimental procedures were performed and national and international therapeutic guidelines were followed. The study was conducted in accordance with the Helsinki Declaration. Clinical case concerns a patient admitted to the hospital and directly managed by the authors.

Patient consent for publication: informed consent was obtained from the patient for the publication of the clinical case.

Availability of data and materials: the clinical case data are available.

Funding: the authors declare they received no financial support for the preparation of this article.

Abstract

This study describes a case of fracture and migration of a peripherally inserted central catheter (PICC) identified as an electrode fragment during an endovascular procedure. It analyzes risk factors, mechanisms, and the role of diagnostic imaging for rapid fragment identification.

A 65-year-old woman with colon carcinoma had a PICC placed for chemotherapy. Routine chest radiography revealed a linear foreign body near the cardiac silhouette. Computed tomography (CT) confirmed a migrated fragment in the right ventricle. Despite no symptoms, urgent intervention was required due to the intracardiac location. The fragment, identified as part of the PICC electrode, was removed percutaneously using a loop snare system without complications, and the patient was discharged in good condition.

Literature indicates that PICC fracture and migration are rare but serious, potentially causing arrhythmias, pulmonary embolism, infections, or valvular damage. Causes include defective materials, mechanical compression (*e.g.*, pinch-off syndrome), improper placement/removal, or material wear. Symptoms vary from none to severe. Early diagnosis relies on radiological surveillance, with chest radiography as the initial step, followed by CT and echocardiography for precise localization. The percutaneous endovascular approach is the preferred treatment due to its efficacy and safety. This case underscores the importance of vigilant radiological follow-up, even in asymptomatic patients, to detect anomalies that could jeopardize safety. CT enabled rapid diagnosis and planning, while the endovascular procedure ensured safe fragment removal and quick recovery. The case highlights the need for healthcare professionals to monitor device integrity and functionality, emphasizing early diagnosis through advanced imaging and prompt intervention to ensure favorable outcomes in complex oncological patients.

Introduction

The peripherally inserted central catheter (PICC) is one of the most common methods of central venous access, particularly useful for patients requiring long-term treatments such as chemotherapy administration, parenteral nutrition, or high-dose antibiotics.¹ The insertion of a PICC is a minimally invasive procedure that offers significant advantages over other types of central venous access, such as central venous catheters (CVC), by reducing the risks of infections and complications associated with invasive surgery.² However, despite its utility, the use of PICCs is not without complications, some of which can be severe, including thrombosis, infection, and, more rarely, catheter fragmentation.³ Catheter fragmentation is a relatively rare but serious complication that occurs when a portion of the catheter detaches and migrates within the vascular system, posing risks of occluding vital blood vessels and causing pulmonary embolism or other thromboembolic events.⁴ This phenomenon can result from multiple factors, including material wear of the catheter, exposure to mechanical forces, or errors during insertion or maintenance of the device.⁵ Although fragmentation is uncommon, its incidence has increased with the advent of thinner and more flexible catheters, which, while offering benefits in terms of comfort and reduced infectious complications, are more susceptible to rupture.⁶ Diagnosing PICC fragmentation can be challenging, as smaller fragments or those located in difficult-to-reach areas are not always visible with traditional imaging techniques.⁷ However, computed tomography (CT) and endovascular ultrasonography have proven useful in identifying and localizing migrated fragments, enabling prompt intervention.⁸ Chest radiography, although the initial imaging modality used to monitor PICCs, may not always be sufficiently sensitive to detect small fragments or confirm their precise location.⁹ Consequently, early diagnosis requires a combination of diagnostic techniques, including CT, magnetic resonance imaging, and ultrasonography, which offer higher accuracy in localizing fragments.¹⁰ The therapeutic approach to PICC fragmentation depends on the location and size of the fragment, as well as the severity of associated complications. In some cases, if the fragments have not caused significant damage, conservative management with regular patient monitoring may be adopted.¹¹ However, when fragments migrate into major vessels such as the superior vena cava and cause obstructions or embolisms, more invasive interventions are necessary, which may include surgical removal or the use of endovascular techniques to extract the fragment less invasively.¹² The endovascular approach, employing techniques such as angioplasty or stent placement, is often preferred due to its minimally invasive nature and lower risk of postoperative complications.¹³ Prevention of PICC fragmentation primarily focuses on selecting more durable materials and proper catheter management. Catheters made from high-quality materials, such as silicone, offer greater flexibility and resistance to fracture compared to polyethylene catheters.¹⁴ Additionally, adequate training of healthcare personnel and adherence to guidelines during catheter insertion and management can significantly reduce the risks of damage and complications.¹⁵ Patients should be educated about signs and symptoms of complications, such as chest pain or respiratory difficulty, which may indicate catheter blockage or fragment release.¹⁶ Therefore, although rare, PICC fragmentation represents a serious complication that can lead to significant clinical events such as pulmonary embolism or central venous thrombosis. Early diagnosis, prompt intervention, and the implementation of preventive strategies are essential to reduce the incidence of this complication and improve clinical outcomes for patients.¹⁷ The use of more robust materials and proper catheter management are critical for patient safety, while advanced diagnostic techniques and endovascular treatments offer promising options for addressing PICC fragmentation.¹⁸

Case Report

A 65-year-old woman with a history of left-sided colon carcinoma at stage III underwent left hemicolectomy, followed by adjuvant chemotherapy with the FOLFOX regimen (5-fluorouracil, leucovorin, and oxaliplatin), in accordance with international guidelines. To ensure safe, continuous, and repeated administration of chemotherapeutic agents, a PICC was placed *via* the left basilic vein, with positioning confirmed through intra-procedural fluoroscopy. The procedure was performed

smoothly, without immediate or delayed complications related to device insertion. The patient completed all scheduled treatment cycles without significant adverse events or the need for premature catheter removal. After completing therapy, the PICC was removed in an outpatient setting, apparently in full and without documented technical difficulties. During follow-up, several weeks later, the patient reported no significant symptoms, particularly no chest pain, dyspnea, palpitations, or signs of infection. However, a routine chest radiograph revealed a hazy, linear opacity projected near the cardiac silhouette, raising suspicion of an intravascular foreign body. The radiological finding prompted clinical suspicion of a possible residual catheter fragment not detected at the time of removal. To clarify the nature and location of the foreign body, a contrast-enhanced chest CT scan was performed, confirming the presence of an approximately 6 cm linear fragment located within the right ventricular cavity, near the apical trabeculation, with no evidence of perforation, pericardial effusion, or associated thrombosis. Despite its intracardiac position, the patient remained completely asymptomatic, with stable vital parameters, normal laboratory tests, and no signs of systemic infection or embolism. Considering the potential risk of delayed complications, such as arrhythmias, distal embolization, or infection of the fragment, a multidisciplinary management approach was initiated, involving interventional radiology, cardiology, and vascular surgery teams. An endovascular percutaneous procedure was scheduled to remove the fragment using a capture system (loop snare) *via* right femoral venous access. The procedure was successfully performed without complications. The patient experienced an uneventful post-procedure course, with complete resolution of the issue and discharge in good general condition (Figure 1).

Discussion

PICC represents one of the most effective solutions for the treatment of patients requiring long-term central venous access without the need for invasive surgical procedures.¹⁹ PICCs are commonly used in clinical settings where central venous access is necessary, such as for the infusion of chemotherapy, long-term antibiotic therapy, and parenteral nutrition.²⁰ Their versatility makes them a popular choice; however, they are not without complications and disadvantages, which must be carefully evaluated by the medical team.²¹

The main advantage of the PICC over other types of central venous access devices, such as CVCs, lies in its ease of insertion and the reduced risk of complications associated with invasive surgical procedures.²² The PICC is inserted *via* a peripheral vein, usually in the forearm, and advanced to the superior vena cava.²³ This percutaneous insertion method significantly reduces the risk of bleeding and infection compared to surgical placement of a CVC, which involves a more invasive procedure.²⁴ Moreover, the PICC can be inserted relatively quickly, even at the patient's bedside, with a lower complication rate compared to other central catheters.²⁵

Another key benefit of the PICC is its potential for long-term use.²⁶ This makes it particularly suitable for oncology patients undergoing prolonged chemotherapy cycles, or for those requiring chronic antibiotic therapy.²⁷ Indeed, numerous studies have demonstrated that the use of PICCs in oncology patients and those with chronic infectious diseases reduces the number of venous punctures and the risk of central venous thrombosis compared to other central devices.^{28,29}

However, prolonged use of the PICC is also associated with specific complications, including infection, occlusion, venous thrombosis, and, more rarely, catheter fragmentation.³⁰ Infectious complications are among the main disadvantages associated with PICC use.³¹ Although infection rates are relatively lower than with CVCs, the risk of local and systemic infections remains, particularly in immunocompromised patients.³² Infections may arise from contamination during catheter insertion, improper handling of the device, or poor maintenance of the insertion site.³³ PICC-related infections can lead to sepsis, a condition that significantly increases patient morbidity and mortality.³⁴

Another associated risk is catheter occlusion.³⁵ Occlusion may occur due to various causes, including blood clots, drug precipitates, or crystallization of infused solutions.³⁶ Management of occlusion may require thrombolytic therapy or, in more severe cases, catheter removal.³⁷

Venous thrombosis is another significant complication.³⁸ Patients with a PICC are at risk of thrombus formation, especially when the catheter remains *in situ* for extended periods.³⁹ PICC-related venous thrombosis can cause pain, swelling, and other clinical manifestations, and may impair blood flow and catheter functionality.⁴⁰ To prevent thrombosis, close patient monitoring and the consideration of prophylactic anticoagulation in high-risk individuals are essential.⁴¹ In some cases, venous thrombosis may progress to more severe complications such as pulmonary embolism, a potentially life-threatening condition.⁴²

Another serious risk associated with PICCs is catheter fragmentation, which occurs when a portion of the catheter breaks off and remains lodged within the patient's vascular system.⁴³ Although PICC fragmentation is rare, it is among the most serious complications, as catheter fragments may migrate to vital organs, such as the lungs or heart, and cause severe damage, including embolism, vascular occlusion, or even myocardial infarction.⁴⁴ Fragmentation can result from mechanical factors, such as repeated catheter movement, improper insertion techniques, or the presence of infection that weakens the catheter material.⁴⁵ The risk of fragmentation is higher in long-term catheters or those subjected to mechanical stress during daily handling.⁴⁶

When fragmentation occurs, immediate management is crucial. One of the first therapeutic measures includes imaging studies, particularly an ultrasound or a CT scan, to locate the detached catheter fragment.⁴⁷ Surgical removal is often necessary and may require a specialized intervention, carrying additional risks for the patient.⁴⁸ Due to the severity of this complication, patients with a PICC should undergo regular monitoring to verify catheter integrity, and any abnormalities, such as sudden pain or difficulty with infusion, must be promptly investigated.⁴⁹ Additionally, patients should be educated about the warning signs of possible catheter fragmentation, such as chest pain or respiratory difficulties, which could indicate embolization of a catheter fragment.⁵⁰

Mechanical complications, such as catheter malpositioning, also represent a significant concern.⁵¹ The PICC must be correctly positioned in the superior vena cava to ensure optimal function.⁵² Incorrect placement can compromise the therapeutic efficacy and increase the risk of complications, such as infection or thrombosis.⁵³ To reduce the risk of malpositioning, catheter placement should be guided and verified through imaging techniques, such as ultrasound or X-ray.⁵⁴

Beyond physical complications, PICC is also associated with psychological considerations for patients.⁵⁵ The placement of a PICC may be perceived as invasive, even though it is less traumatic than other central devices.⁵⁶ Pain management and patient anxiety during and after insertion are important variables that require attention.⁵⁷ Proper evaluation of the patient's tolerability is essential to ensure the PICC can be used without compromising psychological well-being.⁵⁸

It is crucial to highlight the need for adequate training of healthcare professionals managing PICCs.⁵⁹ Insufficient training may lead to errors during catheter insertion, maintenance, or removal, increasing the risk of complications.⁶⁰ Ongoing education and awareness of best practices in PICC management are essential to improve patient outcomes and reduce complication rates.⁶¹ Therefore, PICC represents a safe and effective option for long-term venous access in various clinical contexts.⁶² Although the benefits are numerous, including a lower infection risk and potential for prolonged use, the associated complications, such as infection, occlusion, venous thrombosis, and fragmentation, require careful monitoring and management.⁶³ Careful patient selection, proper staff training, and continuous monitoring are crucial to ensure successful PICC use and to minimize the risks associated with its application.⁶⁴

Conclusions

In conclusion, the incidental detection of PICC fragments represents a clinical complication that, although rare, poses significant potential risks to patient health, including severe outcomes such as pulmonary embolism, sepsis, or damage to surrounding organs. Prompt and accurate identification of such fragments is crucial to prevent fatal complications and to optimize therapeutic intervention.

In this context, diagnostic imaging emerges as an essential element, with advanced techniques such as chest radiography, ultrasonography, and CT playing a fundamental role in localizing and

monitoring the migration of fragments. These non-invasive and highly sensitive diagnostic tools enable precise visualization of fragment position, allowing for timely and effective intervention, which may include surgical removal or other therapeutic maneuvers. The integrated use of imaging technologies also facilitates postoperative monitoring, minimizing the risk of complications and improving the long-term outcome for the patient.

In summary, managing a case of PICC catheter fragmentation requires a multidisciplinary approach, with diagnostic imaging playing a central role in ensuring safe, targeted, and effective treatment.

References

1. Mielke D, Wittig A, Teichgräber U. Peripherally inserted central venous catheter (PICC) in outpatient and inpatient oncological treatment. *Support Care Cancer* 2020;28:4753-60.
2. Adams DZ, Little A, Vinsant C, Khandelwal S. The midline catheter: a clinical review. *J Emerg Med* 2016;51:252-8.
3. Moureau N. Hydrophilic biomaterial intravenous hydrogel catheter for complication reduction in PICC and midline catheters. *Expert Rev Med Devices* 2024;21:207-16.
4. Brescia F, Pittiruti M, Spencer TR, Dawson RB. The SIP protocol update: eight strategies, incorporating Rapid Peripheral Vein Assessment (RaPeVA), to minimize complications associated with peripherally inserted central catheter insertion. *J Vasc Access* 2024;25:5-13.
5. Zochios V, Umar I, Simpson N, Jones N. Peripherally inserted central catheter (PICC)-related thrombosis in critically ill patients. *J Vasc Access* 2014;15:329-37.
6. Moulin V, Monti M. Central venous catheter, PICC-line or Midline: which catheter for my patient? *Rev Med Suisse* 2022;18:121-5. [Article in French].
7. Moureau N, et al. Indications for peripheral, midline and central catheters: summary of the MAGIC recommendations. *Br J Nurs*. 2016. PMID: 27126759.
8. Pinelli F, Muzzi M, Pittiruti M. Should ultrasound evaluation for catheter-related thrombosis always be required before PICC removal? *J Vasc Access* 2024;25:697-702.
9. Santos FKY, Gomes Flumigna RL, Leal Areias L, et al. Peripherally inserted central catheter versus central venous catheter for intravenous access: a protocol for systematic review and meta-analysis. *Medicine* 2020;99:e20352.
10. Lacostena-Pérez ME, Buesa-Escar AM, Gil-Alos AM. Complications related to the insertion and maintenance of peripheral venous access central venous catheter. *Enferm Intensiva* 2019;30:116-26.
11. Rowe MS, Arnold K, Spencer TR. Catheter securement impact on PICC-related CLABSI: a university hospital perspective. *Am J Infect Control* 2020;48:1497-500.
12. Barone G, Natile M, Ancora G. Off label use of PICC as umbilical arterial catheter: clinical case series. *J Vasc Access* 2022;23:796-800.
13. Bing S, Smotherman C, Rodriguez RG, et al. PICC versus midlines: Comparison of peripherally inserted central catheters and midline catheters with respect to incidence of thromboembolic and infectious complications. *Am J Surg* 2022;223:983-7.
14. Chen X, Liang M. A Meta-analysis of incidence of catheter-related bloodstream infection with midline catheters and peripherally inserted central catheters. *J Healthc Eng* 2022;2022:6383777.
15. Tao F, Wang X, Liu J, et al. Perioperative application of midline catheter and PICC in patients with gastrointestinal tumors. *J Buon* 2019;24:2546-52.
16. Yin YX, Gao W, Feng SY, et al. ECG-guided PICC insertion using a new silicon catheter with a conductive tip: a retrospective clinical study. *J Vasc Access* 2022;23:567-73.
17. Yu D, Zhang K, Huang L, et al. Detection of peripherally inserted central catheter (PICC) in chest X-ray images: a multi-task deep learning model. *Comput Methods Programs Biomed* 2020;197:105674.

18. Bredenberg E, Atwater R, Grimm E, et al. Promoting appropriate midline catheter and PICC placement through implementation of an EHR-based clinical decision support tool: an interrupted time-series analysis. *J Hosp Med* 2023;18:483-90.
19. Hu Q, Su Y, Yan L. Effects of peripherally inserted central catheter (PICC) catheterization nursing on bloodstream infection in peripheral central venous catheters in lung cancer: a single-center, retrospective study. *Comput Math Methods Med* 2022;2022:2791464.
20. Cotogni P, Pittiruti M. Focus on peripherally inserted central catheters in critically ill patients. *World J Crit Care Med* 2014;3:80-94.
21. Johansson E, Hammarskjöld F, Lundberg D, Arnlind MH. Advantages and disadvantages of peripherally inserted central venous catheters (PICC) compared to other central venous lines: a systematic review of the literature. *Acta Oncol* 2013;52:886-92.
22. Bahoush G, Salajegheh P, Anari AM, et al. A review of peripherally inserted central catheters and various types of vascular access in very small children and pediatric patients and their potential complications. *J Med Life* 2021;14:298-309.
23. Kim K, Kim Y, Peck KR. Previous peripherally inserted central catheter (PICC) placement as a risk factor for PICC-associated bloodstream infections. *Am J Infect Control* 2020;48:1166-70.
24. Auron M. PICC vs. midline catheter use for short-term indications was linked to major complications at 30 d. *Ann Intern Med* 2022;175:JC47.
25. Caprara J. PICC versus midline. *Home Healthc Now* 2017;35:575-6.
26. Yilmaz FH, Çeri A. A worrisome PICC line complication in a premature newborn: knotted catheter. *Indian J Pediatr* 2021;88:939.
27. Ma S, Shen C, Li Q, et al. Clinical factors of PICC-RVT in cancer patients: a meta-analysis. *Support Care Cancer* 2023;31:393.
28. Van Mechelen K, Mahieu L. A new technique for difficult removal of a peripherally inserted central venous catheter (PICC) in a neonate. *Eur J Pediatr* 2021;180:973-6.
29. Thomsen SL, Boa R, Vinter-Jensen L, Rasmussen BS. Safety and efficacy of midline vs peripherally inserted central catheters among adults receiving IV therapy: a randomized clinical trial. *JAMA Netw Open* 2024;7:e2355716.
30. DeVries M, Sleweon T. Bridging the gap: introduction of an antimicrobial peripherally inserted central catheter (PICC) in response to high PICC central line-associated bloodstream infection incidence. *Br J Nurs* 2021;30:S16-22.
31. Moss JG, Wu O, Bodenham AR, et al. Central venous access devices for the delivery of systemic anticancer therapy (CAVA): a randomised controlled trial. *Lancet* 2021;398:403-15.
32. Ullman AJ, Paterson RS, Schults JA, et al. Do antimicrobial and antithrombogenic peripherally inserted central catheter (PICC) materials prevent catheter complications? An analysis of 42,562 hospitalized medical patients. *Infect Control Hosp Epidemiol* 2022;43:427-34.
33. Johnston AJ, Streater CT, Noorani R, et al. The effect of peripherally inserted central catheter (PICC) valve technology on catheter occlusion rates—the ‘ELeCTRiC’ study. *J Vasc Access* 2012;13:421-5.
34. Sakai H, Iwata M, Terasawa T. External validation of the Michigan PICC catheter-associated bloodstream infections score (MPC score) for predicting the risk of peripherally inserted central catheter-associated bloodstream infections: a single-center study in Japan. *Infect Control Hosp Epidemiol* 2023;44:480-3.
35. Ding Y, Ji L, Hu Y. Effects of tai chi on catheter management and quality of life in tumor patients with PICC at the intermission of chemotherapy: a non-inferiority randomized controlled trial. *Ann Palliat Med* 2020;9:3293-303.
36. Pitiriga V, Bakalis J, Theodoridou K, et al. Lower risk of bloodstream infections for peripherally inserted central catheters compared to central venous catheters in critically ill patients. *Antimicrob Resist Infect Control* 2022;11:137.

37. Joga S, Bansal A, Talwar V, et al. Spontaneous pulmonary arterial embolization: a rare complication of the silicon-based peripherally inserted central venous catheter (PICC). *J Vasc Access* 2022;23:471-3.
38. Narita A, Takehara Y, Maruchi Y, et al. Usefulness of peripherally inserted central catheter port system (PICC-PORT) implantation in the sitting position: a new technique for cases unsuitable for conventional implantation. *Jpn J Radiol* 2023;41:108-13.
39. Taxbro K, Hammarskjöld F, Thelin B, et al. Clinical impact of peripherally inserted central catheters vs implanted port catheters in patients with cancer: an open-label, randomised, two-centre trial. *Br J Anaesth* 2019;122:734-41.
40. Zhang L, Li Y, Zhai Z, Zhang L. The forward withdrawal method for restoration of the PICC catheter ectopic to the internal jugular vein. *Asian J Surg* 2023;46:5728-9.
41. Chopra V, Anand S, Hickner A, et al. Risk of venous thromboembolism associated with peripherally inserted central catheters: a systematic review and meta-analysis. *Lancet* 2013;382:311-25.
42. Furlong-Dillard J, Aljabari S, Hirshberg E. Diagnostic accuracy among trainees to safely confirm peripherally inserted central catheter (PICC) placement using bedside ultrasound. *Br J Nurs* 2020;29:S20-8.
43. Liu X, Tao S, Ji H, et al. Risk factors for peripherally inserted central catheter (PICC)-associated infections in patients receiving chemotherapy and the preventive effect of a self-efficacy intervention program: a randomized controlled trial. *Ann Palliat Med* 2021;10:9398-405.
44. Liu R, Xu H, Pu L, et al. Clinical characteristics of peripherally inserted central catheter-related complications in cancer patients undergoing chemotherapy: a prospective and observational study. *BMC Cancer* 2023;23:894.
45. Erskine B, Bradley P, Joseph T, et al. Comparing the accuracy and complications of peripherally inserted central catheter (PICC) placement using fluoroscopic and the blind pushing technique. *J Med Radiat Sci* 2021;68:349-55.
46. Lawson BT, Zealley IA. Adult 'PICC' device may be used as a tunnelled central venous catheter in children. *Cardiovasc Intervent Radiol* 2018;41:645-52.
47. Fakih M, Sturm L. Paving the PICC journey: building structures, process and engagement to improve outcomes. *BMJ Qual Saf* 2021;30:618-21.
48. Shao G, Zhou X, Zhang S, et al. Cost-utility analysis of centrally inserted totally implanted access port (PORT) vs. peripherally inserted central catheter (PICC) in the oncology chemotherapy. *Front Public Health* 2022;10:942175.
49. Ray-Barruel G, Horowitz J, McLaughlin E, et al. Barriers and facilitators for implementing peripherally inserted central catheter (PICC) appropriateness guidelines: a longitudinal survey study from 34 Michigan hospitals. *PLoS One* 2022;17:e0277302.
50. Zhang J, Si X, Li W, et al. Effect of peripherally inserted central catheter (PICC) parenteral nutrition on immune function and nutritional support after radical gastrectomy for gastric cancer. *Pak J Pharm Sci* 2019;32:1441-45.
51. Bradford NK, Edwards RM, Chan RJ. Normal saline (0.9% sodium chloride) versus heparin intermittent flushing for the prevention of occlusion in long-term central venous catheters in infants and children. *Cochrane Database Syst Rev* 2020;4:CD010996.
52. Schults JA, Kleidon T, Charles K, et al. Peripherally inserted central catheter design and material for reducing catheter failure and complications. *Cochrane Database Syst Rev* 2024;6:CD013366.
53. Burbridge B, Lim H, Dwernychuk L, et al. Comparison of the quality of life of patients with breast or colon cancer with an arm vein port (TIVAD) versus a peripherally inserted central catheter (PICC). *Curr Oncol* 2021;28:1495-506.

54. Tang BZ, Chen CH, Chen H, Ling QY. RCT of reduction in catheter-related complications by using intracavitary electrocardiogram-assisted guidance in neonatal PICC placement. *Sichuan Da Xue Xue Bao Yi Xue Ban* 2021;52:497-502. [Article in Chinese].
55. Braswell LE. Peripherally inserted central catheter placement in infants and children. *Tech Vasc Interv Radiol* 2011;14:204-11.
56. Lee H, Mansouri M, Tajmir S, et al. A Deep-learning system for fully-automated peripherally inserted central catheter (PICC) tip detection. *J Digit Imaging* 2018;31:393-402.
57. Pettit J. Technological advances for PICC placement and management. *Adv Neonatal Care* 2007;7:122-31.
58. Tamura A, Minami K, Tsuda Y, et al. Adolescent eating disorder with catheter-related bloodstream infection. *Pediatr Int* 2021;63:678-84.
59. Yu C, Shulan L, Juan W, et al. The accuracy and safety of using the electrocardiogram positioning technique in localizing the peripherally inserted central catheter tip position: a systematic review and meta-analysis. *Nurs Open* 2022;9:1556-63.
60. Li FX, Su P, Li YP, et al. Definition of catheter jamming: a mechanical analysis of catheter fracture tension and fracture strain. *Technol Health Care* 2022;30:851-7.
61. Mavrovounis G, Mermir M, Chatzis DG, et al. Peripherally inserted central catheter lines for intensive care unit and onco-hematologic patients: a systematic review and meta-analysis. *Heart Lung* 2020;49:922-33.
62. Simonetti G, Sommariva A, Lusignani M, et al. Prospective observational study on the complications and tolerability of a peripherally inserted central catheter (PICC) in neuro-oncological patients. *Support Care Cancer* 2020;28:2789-95.
63. Yu Z, Hu X, Xu X, et al. A nurse-inserted peripherally inserted central catheter program in general pediatrics: a single-center experience. *BMC Pediatr* 2023;23:21.
64. Li J, Hu Z, Luo M, et al. Safety and effectiveness of tunneled peripherally inserted central catheters versus conventional PICC in adult cancer patients. *Eur Radiol* 2024;34:7776-85.




Figure 1. A) A peripherally inserted central catheter with a distal tip located in the right atrium. Thread-like radiopaque image within the cardiac silhouette; B) thread-like hyperdensity within the lumen of the right ventricle; C) venous catheter retrieval performed using a loop snare device; D) electrode fragment removed following endovascular procedure.