

Prospective observational studies in clinical research: pros and cons

Luca Bertolaccini,^{1,2} Matteo Nicoletto,³ Paola Piccolo,⁴ Fulvio Pomero³

¹Department of Oncology and Hemato-Oncology, University of Milan; ²Department of Thoracic Surgery, IEO, European Institute of Oncology IRCCS, Milan; ³Department of Internal Medicine, Ospedale Michele e Pietro Ferrero, Verduno (CN); ⁴Internal Medicine Unit, Ospedale Isola-Tiberina Gemelli Isola, Rome, Italy

Abstract

Prospective observational studies are essential in clinical research, mainly when randomized controlled trials (RCTs) are infeasible, unethical, or impractical. These studies involve the prospective collection of data from cohorts stratified by exposure status, allowing investigators to observe the natural progression of outcomes while preserving temporality. Their main strengths include measuring incidence, assessing multiple outcomes, and avoiding recall bias, thus offering valuable insights into real-world clinical practices. However, these studies also face inherent limitations, most notably the risk of confounding, the absence of randomization, extended duration, and high resource demands. Biases related to selection, performance, and attrition can compromise internal validity if not adequately addressed through rigorous design and statistical adjustment. Although generally favorable, ethical considerations must be carefully navigated, especially concerning data protection and informed consent. While causality cannot be established, prospective observational studies complement RCTs by reflecting effectiveness in typical clinical settings.

This Pros & Cons review examines the structure, advantages, limitations, and methodological considerations of prospective observational studies, incorporating insights from contemporary medical literature and educational frameworks. It also underscores their continued relevance in evidence-based medicine.

Key words: prospective observational study, cohort study, study design, epidemiology, evidence-based medicine, lung cancer.

Correspondence to: Luca Bertolaccini, Department of Oncology and Hemato-Oncology, University of Milan, Via Santa Sofia 9/1 – 20122 Milan, Italy.
Tel: + 39 02 57489665. Fax: +39 02 56562994. E-mail: luca.bertolaccini@unimi.it

Introduction

Observational research, particularly prospective observational studies, is indispensable in clinical epidemiology and healthcare research. These studies allow investigators to explore associations between exposures and outcomes in real-world settings without manipulating variables or randomizing participants. In scenarios where randomized controlled trials (RCTs) may be impractical, unethical, or overly resource-intensive, prospective observational studies offer an ethically sound and pragmatically feasible alternative (Table 1). Moreover, formulating a rigorous and clinically meaningful research question is a fundamental prerequisite for observational research, guiding study design, outcome selection, and statistical interpretation.¹ Prospective observational studies also complement RCTs by providing evidence from routine clinical practice, evaluating heterogeneous patient populations that are frequently underrepresented in experimental trials, and addressing research questions that cannot be investigated ethically or practically through randomization. Consequently, they have become a fundamental component of contemporary evidence-based medicine and real-world evidence generation.^{2,3}

This Pros and Cons explores the strengths and weaknesses of prospective observational studies, incorporating evidence from

the literature and structured teaching material on clinical methodology (Figure 1).

The architecture of a prospective cohort study

In this review, we primarily focus on prospective cohort studies, the most frequently used prospective observational design in clinical research. Participants are enrolled based on exposure status and followed over time to assess the occurrence of predefined outcomes. This temporal framework ensures that the exposure precedes the outcome, an essential feature when inferring causal relationships. Unlike retrospective studies, which analyze pre-existing data, or interventional studies, which actively introduce treatment protocols, prospective designs observe the natural history of events in a controlled yet non-intrusive manner. Temporality is a cornerstone of causal inference, and prospective designs are among the few observational methodologies that permit its assessment effectively.⁴

Observational studies are ideal when ethical or logistical constraints prevent randomized assignment. They are especially well-suited to chronic disease epidemiology, long-term drug safety surveillance, and public health policy evaluation.⁵

Advantages of the design

Prospective observational studies provide several significant advantages. Foremost is their ability to demonstrate temporality, allowing researchers to determine that exposure precedes outcome—an essential, though insufficient, criterion for causality. This is particularly useful in evaluating chronic exposures such as smoking, air pollution, or dietary habits, where experimental manipulation is not feasible.

Unlike RCTs, which frequently enroll highly selected populations under controlled conditions, well-designed prospective cohort studies may better reflect routine clinical practice. However, their external validity depends on the representativeness of the study population and healthcare setting, the completeness of follow-up, and overall data quality. Therefore, generalizability should be interpreted in light of these methodological considerations. As Hess notes, while internal validity may be compromised compared to RCTs, observational studies illustrate effectiveness in practical contexts rather than controlled efficacy.⁵ A typical example of the advantages of a

Table 1. Types of observational studies and their characteristics.

Type	Description	Main characteristics
Cross-sectional study	Assesses exposure and outcome at a single point in time	Quick, low cost; no temporality; useful for prevalence
Case-control study	Compares subjects with an outcome (cases) to those without (controls), looking back at exposures	Efficient for rare diseases; prone to recall and selection bias
Cohort study (prospective)	Follows exposed and unexposed individuals over time to assess outcome occurrence	Good for studying incidence and temporality; time-consuming
Cohort study (retrospective)	Uses historical data to follow groups with different exposures to assess outcome occurrence	Faster and cheaper than prospective cohort; depends on data quality
Ecological study	Analyzes data at the group or population level rather than individual	Useful for population trends; ecological fallacy risk
Nested case-control study	Case-control study within a defined cohort	Reduces selection bias; efficient for biomarker studies




Figure 1. Advantages and disadvantages of prospective observational studies.

prospective observational design is a study conducted at Can Tho Oncology Hospital between 2021 and 2023 that assessed the diagnostic performance of the EU-TIRADS 2017 classification system for malignancy risk stratification.⁶

Moreover, prospective studies mitigate recall bias. Since exposure data are collected before the outcome occurs, they are not influenced by participants' knowledge of their disease status, thereby improving data integrity.⁴ The design also enables the concurrent study of multiple outcomes and risk stratification across subgroups, thereby increasing research efficiency.

Historically, the Framingham Heart Study remains a paradigm for prospective observational research. Initiated in 1948, it identified numerous cardiovascular risk factors, transforming public health policy and clinical guidelines.⁷ Another exemplary initiative is the European Prospective Investigation into Cancer and Nutrition, which has yielded vital insights into the relationships between diet, nutrition, and cancer.⁸

Disadvantages and limitations

Despite their strengths, prospective observational studies are not immune to critical limitations. The most salient disadvantage is their vulnerability to confounding. Because participants are not randomly allocated, differences in baseline characteristics can bias the exposure-outcome relationship. Statistical methods such as multivariable adjustment, propensity score matching, or inverse probability weighting may mitigate, but not eliminate, these concerns.⁵ Residual confounding frequently persists because not all relevant confounders can be measured accurately or even identified. Confounding by indication is particularly important in clinical research, as treatment allocation reflects physicians' judgment and patients' clinical characteristics rather than random assignment.

Another substantial drawback is cost and duration. Longitudinal follow-up is inherently resource-intensive and time-consuming, which may lead to logistical challenges, funding difficulties, and attrition. Loss to follow-up introduces attrition bias, particularly when discontinuation is related to patients' clinical condition, prognosis, or treatment response. Consequently, missing data are often not completely at random and may substantially distort effect estimates. Modern analytical approaches, including multiple imputation and inverse probability of censoring weighting, can partially reduce this bias but rely on specific assumptions.⁴

Additionally, these studies are inefficient for rare outcomes. The need for large cohorts to detect statistically significant effects may render them impractical in settings with low-incidence diseases. Finally, changes in diagnostic criteria, technology, or healthcare delivery over the extended follow-up period may compromise data consistency and introduce misclassification bias. Exposure or outcome misclassification may occur because of changes in diagnostic criteria, evolving technologies, incomplete ascertainment, or measurement error during prolonged follow-up.

Bias and confounding in observational settings

Bias, a systematic deviation from the truth, is a central concern in all observational research. Selection bias, information bias, and confounding are especially problematic in non-randomized designs.⁴ For example, heterogeneity in data collection protocols and clinical expertise can introduce performance bias in extensive multicenter

cohort studies. Moreover, temporal biases may emerge when retrospective elements are incorporated (*e.g.*, using historical data to define baseline exposure).

Confounding occurs when an extraneous variable is associated with the exposure and the outcome, potentially distorting their genuine relationship. For instance, in a study examining the association between exercise and heart disease, dietary habits could be a confounder if healthier diets are more common among those who exercise. Addressing confounding requires careful study design together with appropriate statistical methods, including multivariable regression, propensity score approaches, inverse probability weighting, and sensitivity analyses. Nevertheless, these techniques reduce rather than eliminate confounding, particularly when important variables remain unmeasured.

Methodological enhancements and best practices

To overcome these limitations, a robust methodological framework is essential. This includes the clear articulation of research questions, standardized definitions of exposure and outcomes, and rigorous statistical modeling. Researchers must also prioritize high-quality data collection, ideally supported by electronic health records or prospective registries. In addition, a correct interpretation of absolute and relative effect measures, along with their confidence intervals, is essential for appropriately contextualizing observational findings and their potential clinical implications.⁹ Modern observational research increasingly incorporates causal inference methodologies. Directed acyclic graphs (DAGs) provide a transparent framework for identifying confounders requiring adjustment while avoiding inappropriate adjustment for mediators or colliders. Furthermore, the Target Trial Emulation framework encourages investigators to explicitly define eligibility criteria, treatment strategies, follow-up, outcomes, and analytical approaches before study initiation, thereby reducing common observational biases such as immortal time bias and improving causal interpretation.³

Taken together, this didactic perspective makes it evident that the credibility of a prospective observational study lies in its design and execution. The involvement of experienced statisticians, pre-specified analytical plans, and adherence to reporting guidelines such as the STROBE Statement improve the transparency, completeness, and reproducibility of reporting. However, they do not directly enhance the internal validity of the study itself.^{10,11} Whenever feasible, investigators should pre-specify statistical analysis plans, perform sensitivity analyses to assess the robustness of their findings, and implement appropriate methods for handling missing data.

Ethical and practical considerations

Ethical requirements for prospective observational studies depend on the specific study procedures, informed consent requirements, privacy safeguards, use of routinely collected clinical data, and collection of biological specimens. Although these studies generally involve lower risk than interventional trials, they still require rigorous ethical oversight. Unlike RCTs, they do not withhold treatment or expose participants to untested interventions. Therefore, they are particularly suitable for investigating exposures that would

be unethical to randomize, such as smoking, environmental pollutants, or social determinants of health.

Nonetheless, ethical rigor must still be applied. Informed consent, data confidentiality, and transparency in reporting are mandatory. Moreover, clinicians involved in such studies must maintain clinical equipoise and avoid therapeutic misconception.

Conclusions

Prospective observational studies are potent instruments in the modern research arsenal, offering a compromise between experimental rigor and real-world applicability. While they cannot establish causality with the same authority as randomized trials, they provide critical evidence where RCTs are unfeasible. Their ability to document temporality, monitor long-term outcomes, and generate hypotheses for further investigation makes them invaluable in the clinical research continuum.

However, these advantages come at a cost—the risk of bias, confounding, and logistical complexity. By adhering to methodological best practices and maintaining a critical perspective, researchers can harness the full potential of prospective observational studies while recognizing their limitations. As the boundaries of clinical inquiry continue to expand, carefully designed observational research will only grow more central in guiding patient care and public health policy.

References

1. Bonaventura A, Buso R, Dentali F. The research question: a first, essential step in study design. *Ital J Med* 2026;20:2130.
2. Concato J, Shah N, Horwitz RI. Randomized, controlled trials, observational studies, and the hierarchy of research designs. *N Engl J Med* 2000;342:1887-92.
3. Hernan MA, Robins JM. Using big data to emulate a target trial when a randomized trial is not available. *Am J Epidemiol* 2016; 183:758-64.
4. Thiese MS. Observational and interventional study design types; an overview. *Biochem Med* 2014;24:199-210.
5. Hess DR. Observational studies. *Respir Care* 2023;68:1585-97.
6. Van Kha V, Thi Huong Ly T, Thi Thanh Hoa P, Huynh Nhu V. Employing the European Thyroid Imaging Reporting and Data System 2017 classification in a malignancy risk stratification system for thyroid nodules at Can Tho Oncology Hospital from 2021 to 2023. *Ital J Med* 2025;19:1852.
7. Kannel WB, Dawber TR, Kagan A, et al. Factors of risk in the development of coronary heart disease—six year follow-up experience. The Framingham Study. *Ann Intern Med* 1961;55:33-50.
8. Riboli E, Hunt KJ, Slimani N, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr* 2002;5:1113-24.
9. Maino A, Bertù L, Squizzato A, et al. Quantifying treatment effect: relative and absolute measures in clinical research. *Ital J Med* 2026;20:2386.
10. Vandenbroucke JP, von Elm E, Altman DG, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *Ann Intern Med* 2007;147:W163-94.
11. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ* 2007;335: 806-8.

Received: 13 May 2026; Accepted: 26 July 2026; Early view: 28 August 2026.

Contributions: Luca Bertolaccini: study concept, manuscript original drafting; Fulvio Pomero: manuscript writing and editing. All authors have read and approved the final version of the manuscript and agree to be accountable for all aspects of the work. Conflict of interest: the authors declare that they have no competing interests.

Ethics approval and consent to participate: not applicable.

Informed consent: not applicable.

Patient consent for publication: not applicable.

Availability of data and materials: not applicable.

Acknowledgments: this work was partially supported by the *Ministero della Salute* with *Ricerca Corrente* and *5x1000* funds.

Declaration of generative artificial intelligence and artificial intelligence-assisted technologies in the writing process: no AI used in the writing process.

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).